

Systemic Inflammation Mediates the Effects of Endotoxemia in the Mechanisms of Heat Stroke

Chin Leong Lim¹ and Katsuhiko Suzuki^{2*}

¹Lee Kong Chian School of Medicine, Nanyang Technological University, 59 Nanyang Drive, Singapore

²Faculty of Sport Sciences, Waseda University, Mikajima, Tokorozawa, Saitama, Japan

*Corresponding author: Katsuhiko Suzuki, MD, PhD, Faculty of Sport Sciences, Waseda University, 2-579-15 Mikajima, Tokorozawa, Saitama 359-1192, Japan, Tel & Fax: +81-4-2947-6898; E-mail: katsu.suzu@waseda.jp

Received date: November 30, 2016; Accepted date: December 24, 2016; Published date: December 31, 2016

Copyright: © 2016 Lim CL and Suzuki K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Heat stroke is triggered by heat, but is driven by endotoxemia and the downstream effects of systemic inflammation, acute phase response and the pyrogenic response. Whereas heat stroke and its related fatality commonly occur at core temperature (T_c) >40°C, healthy individuals have tolerated T_c 40°C-42°C without symptoms of heat stroke, suggesting that besides hyperthermia, there are other factors that can also cause heat stroke. The “dual-pathway model (DPM)” suggests that heat stroke is triggered by the endotoxemia pathway at T_c up to ~42°C and independently, by heat toxicity at T_c >42°C. The second pathway in the DPM is based on evidence showing that cytoskeletal structures start to disintegrate at ambient temperature >41.5°C. Since most exertional stroke cases occur at T_c <42°C, the endotoxemia pathway, and not heat, might be the primary cause of heat stroke in the active population. Current evidence suggests that exercising under a poor state of health and a compromised immune system may also cause heat stroke, independent from the effects of hyperthermia. Strategies to prevent heat stroke should put equal emphasis on maintaining a good state of health and immune function. The current practice of focusing primarily on heat strain and hydration to prevent heat stroke may not have comprehensively addressed the pathophysiology of heat stroke and may explain why heat stroke continues to occur after more than 2000 years.

Keywords: Heat stress; Exercise; Endotoxemia; Leakage of gram-negative bacteria (bacterial translocation); Lipopolysaccharides (LPS); Gut barrier; Inflammatory cytokines

Introduction

Heat stroke is the fatal form of heat injury that is often presented with hyperthermia and symptoms of central nervous system (CNS) dysfunction and can lead to coma and death [1,2]. Heat stroke is also an ancient illness dating back more than two thousand years and its pathology has been attributed to the effects of hyperthermia and heat toxicity [3-5]. The roles of hyperthermia and heat toxicity in causing heat stroke have not been proven empirically, but form the basis for the current understanding, prevention and treatment of heat stroke [5,6]. However, evidence presented in the last two decades suggests that heat- and exercise-induced endotoxemia and the downstream consequences of systemic inflammation may play important roles in triggering and driving the mechanisms of heat stroke [2,7-9]. Therefore, this short review aims to summarize and discuss the evidence supporting the roles of endotoxemia and systemic inflammation in triggering and driving the mechanisms of heat stroke, independent from the effects of heat.

“Dual-Pathway Model (DPM)” of Heat Stroke

The earlier proponents of endotoxemia in the mechanisms of heat stroke advocated that heat stroke is triggered by heat but is driven by endotoxemia and the downstream effects of systemic inflammation, acute phase response and the pyrogenic response [2,9,10]. However, whereas heat stroke and its related fatality commonly occur at core

temperature (T_c) >40°C [11], healthy individuals have tolerated T_c 40°C-42°C without symptoms of heat stroke [12,13]. This evidence suggests that besides hyperthermia, there are other factors that can also cause heat stroke, which supports the notion that heat and endotoxemia may activate the mechanisms of heat stroke through independent pathways [8].

The “dual-pathway model (DPM)” of heat stroke suggests that heat stroke is triggered by the endotoxemia pathway at T_c up to ~42°C and independently, by heat toxicity at T_c >42°C [8]. The second pathway in the DPM is based on *in vitro* evidence showing that cytoskeletal structures start to disintegrate at ambient temperature >41.5°C [14]. Since most exertional stroke cases occur at T_c <42°C [8,11], the endotoxemia pathway, and not heat, might be the primary cause of heat stroke in the active population.

Gut Barrier Response to Exercise and Heat Stresses

Proponents of the endotoxemia pathway agree that the pathophysiology of heat stroke originates from the gut, which houses hundreds of millions of bacteria [9,15]. The gut epithelium functions as a physical barrier that separates the septic condition in the gut space from the aseptic condition in circulating blood. However, the permeability of the gut-epithelium is increased during intense exercise and heat stress exposure, which promotes the leakage of gram-negative bacteria (endotoxins, a.k.a. lipopolysaccharides, LPS) from the gut space into the portal circulation [16,17]. From the portal circulation, LPS is transported to the liver, where LPS is scavenged and removed from the body by Kupffer cells (hepatic macrophages) [9,10]. During prolonged intense exercise, LPS influx may overwhelm LPS clearance in the liver [9,15,18], leading to the leakage of LPS from the liver into

the central circulation, where LPS can be removed by high density lipoprotein, LPS-specific antibodies and the innate immune system [19].

Pro-inflammatory Cytokine Response to Endotoxemia

A sub-lethal increase in the circulating concentration of LPS can induce the production of pro-inflammatory cytokines, such as tumor necrosis factor (TNF)- α , interleukin (IL)-1 β , and IL-6, which have both inflammatory and pyrogenic functions [2,10]. In a healthy state, the anti-LPS mechanisms can scavenge and remove LPS from the body through the reticuloendothelial system in the liver [8]. However, under conditions of immune suppression, LPS concentration in the blood can increase to the threshold of triggering the systemic inflammatory response [8]. The downstream effects of this “shock” response include disseminated intravascular coagulation (DIC), multi-organ failure (MOF), haemorrhage and CNS disturbances, which are commonly reported in heat stroke victims [20-22]. Therefore, the endotoxemia pathway implies that during exposure to heat and exercise stresses, the integrity of the gut barrier may play important “gatekeeping” roles in blocking the cascade of events that can lead to the occurrence of heat stroke. Under conditions where the immune system is compromised and is unable to cope with the influx of LPS from the gut, endotoxemia can progress to activate the mechanisms of heat stroke through the downstream effects of septicaemia and systemic inflammation.

Evidence Derived from Studies on Healthy Athletes

The roles of immune disturbances as an independent trigger of heat stroke are supported by evidence derived from healthy individuals, heat stroke victims and from animals that were exposed to lethal heat stress. Healthy runners have increased blood concentration of LPS during endurance races [23-25] and gut permeability changes during exercise were positively associated with exercise intensity [26]. Moreover, the increase in gut permeability during intense exercise resulted in endotoxemia only when running in warm, but not in cool conditions [15] and aerobic training can modulate the increase in circulating concentration of LPS during exercise in the heat [27]. However, four weeks of daily probiotic supplementation (45 billion CFU) did not have significant effects on serum LPS concentration and gut permeability during an acute bout of intense exercise in the heat [28]. This evidence demonstrates the occurrence of mild endotoxemia and the associated increase in gut permeability during intense exercise and supports the possible roles of gut-related LPS translocation in the mechanisms of heat stroke.

Evidence Derived from Heat Stroke Victims and Animal Studies

In heat stroke victims, blood samples collected at the emergency department had increased concentrations of endotoxin and pro-inflammatory cytokines [29,30]. These immune changes are consistent with common pathological findings in deceased heat stroke victims, which include systemic inflammation, DIC, MOF, necrosis, hemorrhage, and symptoms of CNS failure [19,20,31,32]. Except for hyperthermia, these clinical symptoms in heat stroke victims are similar to those reported in patients with sepsis (endotoxemia) [9]. Epidemiological evidence from a military hospital in Thailand also supports the notion that exercising with a suboptimal immune system can increase the risk of having heat stroke [33]. Twelve years of

medical records showed that more than 95% of heat stroke patients treated in the military hospital had mild fever, upper respiratory tract infection or diarrhea prior to the heat stroke event [33]. This evidence further supports the central role of a compromised immune system in triggering the mechanisms of heat stroke, possibly through endotoxemia-induced systemic inflammation. In animal studies, inhibiting endotoxemia through gut cleansing and injection of steroids, anti-LPS antibody or antibiotic drugs protected 100% of dogs [34], rabbits [35], monkeys [36-38] and rats [39] from lethal heat stress, when almost all animals in the control group died under the same conditions. Taken together, the evidence presented suggests that the occurrence of endotoxemia or having an infection before and during prolonged intense exercise and heat exposure may progress to trigger the systemic inflammatory response, which may function as the switch that triggers or inhibits the mechanisms of heat stroke. Protecting the health of the individuals, and the immune system in particular, during prolonged periods of intense physical training and heat exposure may play critical roles in protecting against heat stroke due to the endotoxemia pathway.

Conclusion and Future Directions

The current evidence suggests that exercising under a poor state of health and a compromised immune system may cause heat stroke, independent from the effects of hyperthermia. The evidence also suggests that healthy active individuals can tolerate a higher degree of heat stress (up to $T_c \sim 42^\circ\text{C}$) without triggering heat stroke. Strategies to prevent heat stroke, especially in the active population, should put equal emphasis on maintaining a good state of health and immune function. The current practice of focusing primarily on controlling heat strain and on promoting hydration to prevent heat stroke may not address the pathophysiology of heat stroke comprehensively [7] and may explain why heat stroke continues to occur after more than 2000 years. Although preventing hyperthermia [40] and ensuring fluid ingestion [41,42] can attenuate the systemic inflammatory response during exhaustive exercise [22,43-46], the other countermeasures to improve general and specific health conditions, such as functional food [45-50] to promote gut barrier protection and the appropriate immune surveillance, should be examined in the future studies.

Acknowledgements

This work was partly supported by a Grant-in-Aid for the Scientific Research (A) and Strategic Research Foundation at Private Universities from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

1. Ash CJ (2001) Heat stroke. *Mil Med* 166: 5.
2. Bouchama A, Knochel JP (2002) Heat Stroke. *N Engl J Med* 346: 1978-1988.
3. Bouchama A (1995) Heatstroke: a new look at an ancient disease. *Intensive Care Med* 21: 623-625.
4. Eichler AC, McFee AS, Root HD (1969) Heat stroke. *Am J Surgery* 118: 855-863.
5. Roberts WO, Dorman JC, Bergeron MF (2016) Recurrent heat stroke in a runner: Race simulation test for return to activity. *Med Sci Sports Exerc* 48: 785-789.

6. American College of Sports Medicine (1987) Position stand: The prevention of thermal injuries during distance running. *Med Sci Sports Exerc* 19: 529-533.
7. Lim CL (2016) Look beyond thermoregulation and hydration in the diagnosis of heat stroke. *Med Sci Sports Exerc* 48: 2583.
8. Lim CL, Mackinnon LT (2006) The role of exercise-induced immune system disturbances in the pathology of heat stroke: The dual pathway model of heat stroke. *Sports Med* 36: 39-64.
9. Moseley P, Gisolfi CV (1993) New frontiers in thermoregulation and exercise. *Sports Med* 16: 163-167.
10. Hales RJS, Sakurada S (1998) Heat tolerance: A role for fever? *Ann NY Acad Sci* 856: 188-205.
11. Shapiro Y, Seidman DS (1990) Field and clinical observations of exertional heat stroke patients. *Med Sci Sports Exerc* 22: 6-14.
12. Byrne C, Lee J, Chew SAN, Lim CL, Tan YM (2006) Continuous thermoregulatory responses to mass-participation distance running in heat. *Med Sci Sports Exerc* 38: 803-810.
13. Pugh LGCE, Corbett JL, Johnson RH (1967) Rectal temperatures, weight losses and sweat rates in marathon running. *J Appl Physiol* 23: 347-352.
14. Brooks GA, Fahey TD, White TP (1996) Exercise physiology: Human bioenergetics and its applications. Mayfield publishing company, California. p. 504.
15. Gisolfi CV, Mora F (2000) The hot brain: survival, temperature and the human body. MIT Press, Massachusetts, pp. 1-13, 94-119, 157-163, 171-174, 191-215.
16. Lambert GP, Gisolfi CV, Berg DJ, Moseley PL, Oberley LW, et al. (2002) Hyperthermia-induced intestinal permeability and role of oxidative and nitrosative stress. *J Appl Physiol* 92: 1750-1761.
17. Yeh YJ, Law LYL, Lim CL (2013) Gastrointestinal response and endotoxemia during intense exercise in hot and cool environments. *Eur J Appl Physiol* 113: 1575-1583.
18. Dubose DA, McCreary J, Sowders L, Goode L (1983) Relationship between rat heat stress mortality and alterations in reticuloendothelial carbon clearance function. *Aviat Space Environ Med* 54: 1090-1095.
19. Doran JE (1992) Biological effects of endotoxin. In: Gut-derived infectious toxic shock Current studies in hematology and blood transfusion. Karger, Basel. p. 66-99.
20. Charan NB, Robinson WA, Mathew M (1975) Heat stroke, disseminated intravascular coagulation and death in a long distance runner. *J Assoc Physicians India* 23: 917-919.
21. Shibolet S, Coll R, Gilat T, Sohar E (1967) Heat stroke: its clinical picture and mechanism in 36 cases. *Q J Med* 36: 525-547.
22. Suzuki K, Nakaji S, Yamada M, Totsuka M, Sato K, et al. (2002) Systemic inflammatory response to exhaustive exercise: Cytokine kinetics. *Exerc Immunol Rev* 8: 6-48.
23. Camus G, Poortman JR, Nys M, Deby-Dupont G, Duchateau J, et al. (1997) Mild endotoxemia and the inflammatory response induced by a marathon race. *Clin Sci* 92: 415-422.
24. Jeukendrup AE, Vet-Joop K, Sturk A, Stegen JHJC, Senden J, et al. (2000) Relationship between gastro-intestinal complaints and endotoxemia, cytokine release and the acute-phase reaction during and after a long-distance triathlon in highly trained men. *Clin Sci* 98: 47-55.
25. Ng QY, Lee KW, Byrne C, Ho TF, Lim CL (2008) Plasma endotoxin and immune responses during a 21-km road race under a warm and humid environment. *Ann Acad Med Singapore* 37: 307-314.
26. Pals KL, Chang RT, Ryan AL, Gisolfi CV (1997) Effect of running intensity on intestinal permeability. *J Appl Physiol* 82: 571-576.
27. Selkirk GA, McLellan TM, Wright HE, Rhind SG (2008) Mild endotoxemia, NF- κ B translocation, and cytokine increase during exertional heat stress in trained and untrained individuals. *Am J Physiol* 295: R611-623.
28. Shing CM, Peake J, Lim CL, Briskey D, Walsh NP, et al. (2014) Effects of probiotics supplementation on gastrointestinal permeability, inflammation and exercise performance in the heat. *Eur J Appl Physiol* 114: 93-103.
29. Bouchama A, Al-Sedairy S, Siddiqui S, Shail E, Razeig M (1993) Elevated pyrogenic cytokines in heatstroke. *Chest* 104: 1498-1502.
30. Bouchama A, Parhar S, El-Yagi A, Sheth K, Al-Sedairy S (1991) Endotoxemia and the release of tumor necrosis factor and interleukin-1-alpha in acute heat stroke. *J Appl Physiol* 70: 2640-2644.
31. Chao TC, Sinniah R, Pakiam JE (1981) Acute heat stroke deaths. *Pathology* 13: 145-156.
32. Malamud N, Haymaker W, Custer R (1946) Heat stroke: a clinico-pathologic study of 125 fatal cases. *Mil Surg* 99: 397-449.
33. Sithinamsuwan P, Piyavechvirathanat K, Kitthaweesin T, Chusri W, Orrawanhanonthai P, et al. (2009) Exertional heat stroke: Early recognition and outcome with aggressive combined-cooling - a 12-year experience. *Mil Med* 174: 496-502.
34. Bynum GD, Brown J, Dubose DA, Marsili M, Leav I, et al. (1979) Increased survival in experimental dog heatstroke after reduction of gut flora. *Aviat Space Environ Med* 50: 816-819.
35. Butkow N, Mitchell D, Laburn H, Kenedi E (1984) Heat stroke and endotoxemia in rabbits. In: Hales JRS (ed.). *Thermal Physiology*. Raven Press, New York. pp: 511-514.
36. Gathiram P, Wells MT, Brock Utne JG, Gaffin SL (1987) Antilipopolysaccharide improves survival in primates subjected to heat stroke. *Circ Shock* 23: 157-164.
37. Gathiram P, Wells MT, Brock UJG, Gaffin SL (1988) Prophylactic corticosteroid increases survival in experimental heat stroke in primates. *Aviat Space Environ Med* 59: 352-355.
38. Gathiram P, Wells MT, Brock UJG, Wessels BC, Gaffin SL (1987) Prevention of endotoxaemia by non-absorbable antibiotics in heat stress. *J Clin Pathol* 40: 1364-1368.
39. Lim CL, Wilson G, Brown L, Coombes JS, Mackinnon LT (2007) Pre-existing inflammatory state compromises heat tolerance in rats exposed to heat stress. *Am J Physiol* 292: R186-R194.
40. Kondou T, Ogawa K, Terada O, Kim K, Okutsu M, et al. (2009) Cytokine response to exercise during exposure to cold. *Jap J Complement Altern Med* 6: 89-95.
41. Suzuki K, Hashimoto H, Oh T, Ishijima T, Mitsuda H, et al. (2013) The effects of sports drink osmolality on fluid intake and immunoendocrine responses to cycling in hot conditions. *J Nutr Sci Vitaminol* 59: 206-212.
42. Suzuki K, Shiraishi K, Yoshitani K, Sugama K, Kometani T (2014) The effect of a sports drink based on highly branched cyclic dextrin on cytokine responses to exhaustive endurance exercise. *J Sports Med Phys Fitness* 54: 622-630.
43. Suzuki K, Nakaji S, Yamada M, Liu Q, Kurakake S, et al. (2003) Impact of a competitive marathon race on systemic cytokine and neutrophil responses. *Med Sci Sports Exerc* 35: 348-355.
44. Suzuki K, Totsuka M, Nakaji S, Yamada M, Kudoh S, et al. (1999) Endurance exercise causes interaction among stress hormones, cytokines, neutrophil dynamics, and muscle damage. *J Appl Physiol* 87: 1360-1367.
45. Hashimoto H, Ishijima T, Suzuki K, Higuchi M (2016) The effect of the menstrual cycle and water consumption on physiological responses during prolonged exercise at moderate intensity in hot conditions. *J Sports Med Phys Fitness* 56: 951-960.
46. Hashimoto H, Ishijima T, Hayashida H, Suzuki K, Higuchi M (2014) Menstrual cycle phase and carbohydrate ingestion alter immune response following endurance exercise and high intensity time trial performance test under hot conditions. *J Int Soc Sports Nutr* 11: 39.
47. Lim CL, Pyne D, Horn P, Kalz A, Saunders P, et al. (2009) The effects of increased endurance training load on biomarkers of heat intolerance during intense exercise in the heat. *Appl Physiol Nutr Metab* 34: 616-624.
48. Peake J, Peiffer JJ, Abbiss CR, Nosaka K, Laursen PB, et al. (2008) Carbohydrate gel ingestion and immunoendocrine responses to cycling in temperate and hot conditions. *Int J Sport Nutr Exerc Metab* 18: 229-246.
49. Peake JM, Suzuki K, Coombes JS (2007) The influence of antioxidant supplementation on markers of inflammation and the relationship to oxidative stress after exercise. *J Nutri Biochem* 18: 357-371.

50. Suzuki K, Takahashi M, Li CY, Lin SP, Tomari M, et al. (2015) The acute effects of green tea and carbohydrate co-ingestion on systemic inflammation and oxidative stress during sprint cycling. Appl Physiol Nutr Metabol 40: 997-1003.